



**Prepn. of (pyrimid-2-yl-thio- or seleno-) acetic acid derivs. - by reacting the corresp. chloro:alkanoyl-amino cpd. with a rhodanide and water or an alcohol**

**Patent number:** DE4119767  
**Publication date:** 1992-12-17  
**Inventor:** GEWALD KARL PROF DR (DE); SCHAEFER HARRY DR SC NAT (DE); JESCHKE TORSTEN DIPL CHEM (DE); ECKERT KATRIN DIPL CHEM (DE); FAUST GOTTFRIED DR (DE); LABAN GUNTER DR (DE)  
**Applicant:** DRESDEN ARZNEIMITTEL (DE)  
**Classification:**  
**- international:** C07D239/88; C07D239/94; C07D471/04; C07D487/04; C07D495/04; C07D513/04  
**- european:** C07D239/95; C07D487/04; C07D495/04; C07D513/04  
**Application number:** DE19914119767 19910615  
**Priority number(s):** DE19914119767 19910615

**Abstract of DE4119767**

Prepn. of (pyrimid-2-ylthio or seleno)-ethanoic acid derivs. of formula (I) comprises reacting a cpd. of formula (II) with a cpd. of formula MXCN (III) and with a cpd. of formula HR<sup>2</sup> (IV) with heating. In formulae, R<sup>1</sup> is amino or OH; R<sup>2</sup> is OH or 1-4C alkoxy; X is S or Se; A is a substd. aromatic ring condensed in the 4,5-position of the pyrimidine ring and opt. mono- or di-substd. with NO<sub>2</sub>, halo, Me or MeO, or A is a 5-7 membered heterocyclic ring, condensed in the 4,5-position of the pyridimine ring and opt. mono- or di-substd. with Me, methylamino, dimethylamino, MeS, ethanoyl, allyl, 1-4C alkoxy carbonyl tetramethylene, Ph, anilino or anilinocarbonyl, (these last 3 gps. are opt. substd. with halogen and/or MeO); Y is nitrile or 1-4C alkoxy carbonyl; Hal is halo, pref. Cl or Br; and M is NH<sub>4</sub>, Na or K. USE/ADVANTAGE - (I) are intermediates in organic syntheses, e.g., in the prepn. of pharmaceuticals. Some (I) are themselves pharmaceuticals.



Data supplied from the esp@cenet database - Worldwide